

WHAT IS CLAIMED IS:

1. A high-throughput method of distinguishing at least one molecule individually in a sample comprising multiple molecules, which method comprises:
 - (i) subjecting a sample comprising multiple molecules, at least one molecule of which is detectably labeled, to electrophoresis,
 - (ii) imaging the electrophoretic mobility of each detectably labeled molecule over time by detecting the position of the detectable label of each detectably labeled molecule over time and, optionally, at the same time, dispersing the imaging by a transmission grating for spectroscopic analysis, and
 - (iii) determining the electrophoretic mobility of each detectably labeled molecule and, optionally, determining the molecular spectrum of each detectably labeled molecule,thereby distinguishing at least one molecule individually in a sample comprising multiple molecules.
2. The high-throughput method of claim 1, wherein said multiple molecules in said sample are not amplified prior to being subjected to electrophoresis.
3. The high-throughput method of claim 1, wherein said at least one molecule is a nucleic acid.
4. The high-throughput method of claim 3, wherein said nucleic acid is detectably labeled with a fluorescent label.
5. The high-throughput method of claim 1, wherein said at least one molecule is a protein.
6. The high-throughput method of claim 5, wherein said protein is detectably labeled with a fluorescent label.
7. The high-throughput method of claim 1, wherein said at least one molecule is a small molecule.

8. The high-throughput method of any of claims 1-7, wherein said sample comprises a buffer.
9. The high-throughput method of claim 8, wherein said buffer is photobleached.
10. The high-throughput method of claim 8, wherein said buffer comprises a sieving matrix.
11. The high-throughput method of claim 4 or 6, wherein said fluorescent label is induced to fluoresce by a laser.
12. The high-throughput method of any of claims 4, 6 and 11, wherein fluorescence from said fluorescent label is focused on an imaging means.
13. The high-throughput method of claim 12, wherein said imaging means is an intensified CCD camera.
14. The high-throughput method of any of claims 11-13, wherein said laser generates extraneous light and said extraneous light is eliminated through the use of an equilateral prism and at least one optical pinhole positioned before said imaging means.
15. The high-throughput method of claim 12 or 13, wherein one or more optical filters are positioned in front of said imaging means.
16. The high-throughput method of any of claims 1-15, wherein said electrophoretic mobility is measured by a method selected from the group consisting of the multiframe method, the streak method and the multispot method.
17. The high-throughput method of any of claims 1-16, wherein said electrophoretic mobility is imaged in less than about 5 milliseconds.

23. The system of claim 21 or 22, wherein said laser generates extraneous light and said system further comprises an equilateral prism and at least one optical pinhole before said imaging means, wherein said equilateral prism and said at least one optical pinhole eliminate said extraneous light prior to it impinging on said imaging means.

24. The system of any of claims 21-23, wherein said imaging means is an intensified CCD camera.

25. The system of claim 24, which further comprises a microscope objective between said electrophoretic sample channel and said imaging means, wherein said microscope objective focuses the fluorescence from said fluorescent label onto said imaging means.

26. The system of claim 21 or 22, which further comprises one or more optical filters positioned in front of said imaging means.

27. The system of any of claims 21-26, wherein said imaging means images the electrophoretic mobility of a detectably labeled molecule in said sample in less than about 5 milliseconds.

28. The system of any of claims 21-26, wherein said imaging means images the electrophoretic mobility of at least about 200 detectably labeled molecules every 10 milliseconds.

29. The system of any of claim 21-26, wherein said imaging means images the electrophoretic mobility of at least about 2,500 detectably labeled molecules every 25 milliseconds.

30. A high-throughput method of distinguishing at least one molecule individually in a sample comprising multiple molecules, which method comprises:

- (i) introducing a sample comprising multiple molecules in free solution, at least one molecule of which is detectably labeled, into a sample channel,
- (ii) simultaneously imaging the position of each detectably labeled molecule by detecting the position of the detectable label of each detectably labeled molecule and dispersing the imaging by a transmission grating for spectroscopic analysis, and

(iii) determining the molecular spectrum of each detectably labeled molecule,

thereby distinguishing at least one molecule individually in a sample comprising multiple molecules.

31. The high-throughput method of claim 30, wherein said multiple molecules in said sample are not amplified prior to being introduced into said sample channel.

32. The high-throughput method of claim 30, wherein said at least one molecule is a nucleic acid.

33. The high-throughput method of claim 32, wherein said nucleic acid is detectably labeled with a fluorescent label.

34. The high-throughput method of claim 30, wherein said at least one molecule is a protein.

35. The high-throughput method of claim 34, wherein said protein is detectably labeled with a fluorescent label.

36. The high-throughput method of claim 30, wherein said at least one molecule is a small molecule.

37. The high-throughput method of any of claims 29-36, wherein said sample comprises a buffer.

38. The high-throughput method of claim 37, wherein said buffer is photobleached.

39. The high-throughput method of claim 33 or 35, wherein said fluorescent label is induced to fluoresce by a laser.

48. A system for use in the method of claim 30, which system comprises:
- (i) a sample channel, into which is introduced a sample comprising multiple molecules in free solution, at least one molecule of which is detectably labeled with a fluorescent label,
 - (ii) a light source comprising or consisting essentially of at least one wavelength of light that causes at least one molecule in said sample comprising multiple

molecules that is detectably labeled with a fluorescent label to fluoresce, wherein said light source irradiates said sample channel,

(iii) an imaging means, wherein said imaging means images the position of each detectably labeled molecule in said sample, and,

(iv) a transmission grating, which simultaneously disperses the imaging of the position of each detectably labeled molecule in said sample.

49. The system of claim 48, which further comprises a lens between said light source and said sample channel, wherein said lens focuses said light at normal incidence to said sample channel.

50. The system of claim 49, wherein said laser generates extraneous light and said system further comprises an equilateral prism and at least one optical pinhole before said imaging means, wherein said equilateral prism and said at least one optical pinhole eliminate said extraneous light prior to it impinging on said imaging means.

51. The system of any of claims 48-50, wherein said imaging means is an intensified CCD camera.

52. The system of claim 51, which further comprises a microscope objective between said sample channel and said imaging means, wherein said microscope objective focuses the fluorescence from said fluorescent label onto said imaging means.

53. The system of any of claims 48-52, which further comprises one or more optical filters positioned in front of said imaging means.

54. The system of any of claims 48-53, wherein said imaging means images the position of a detectably labeled molecule in said sample in less than about 0.05 milliseconds.

55. The system of any of claims 48-54, wherein said imaging means images the position of at least about 200 detectably labeled molecules every 0.10 milliseconds.

56. The system of any of claims 48-54, wherein said imaging means images the position of at least about 2,500 detectably labeled molecules every 0.25 milliseconds.